

STOMA PRESCRIBING GUIDELINES		
Туре	Average wear time	Average quantity per month
ILEOSTOMY		
One-piece	Pouch changed every 1 - 3 days	10 - 30 pouches
drainable pouch		
Two-piece	Baseplate changed 2 - 3 times	8 - 12 baseplates
drainable pouch	a week	10 - 20 may share
	Pouch changed every 1 - 3 days	10 - 30 pouches
COLOSTOMY		
One-piece pouches closed pouch	Pouch changed 1 - 3 times a day	30 - 90 pouches
Two-piece baseplates closed pouch	Baseplate changed 2 - 3 times a week	8 - 12 baseplates
	Pouch changed 1 - 3 times a day	30 - 90 pouches
variety of one and two piece appliances an UROSTOMY One-piece urostomy pouch Two-piece urostomy pouch	nd additional products used and a wide variety of c Pouch changed every 1 - 3 days Baseplate changed 2 - 3 times a week	10 - 30 pouches 8 - 12 baseplates
	Pouch changed every 1 - 3 days	10 - 30 pouches
Open night drainage bag	Bag changed weekly	4 bags
Single use drainage bag	Bag changed daily	30 bags
CONVEX PRODUCTS	Convex products should only be used on the recommendation of the Stoma Care Nurse.	
SUPPLEMENTARY ITEMS	Additional items may be essential for problem solving with stomal complications or to extend wear time and increase confidence. These include barrier cream, skin protector wipes/spray, powders, adhesive remover, washers and pastes, deodorants, retention strips and support garments.	
The quantities listed above are guidelines only. Some patients may require a higher or lower quantity than stated. If there are any prescribing concerns please refer to the patient's Stoma Care Nurse for details or an assessment.		

Anyone wishing to use this material is requested to acknowledge the source: "PIPs Forum, Stoma prescribing guidelines".

Incorporating Patient, Professional and Industry Groups concerned with stoma and continence care