

AZATHIOPRINE SHARED CARE PROTOCOL

This Shared Care Protocol is for patients initiated on Azathioprine under the Rheumatology Department. The consultant will have detailed the expected treatment regimen in the clinic letter. The first 3 months of prescribing and monitoring will be undertaken by the Rheumatology department. Updates of dosages and results will be completed by the specialist nurses and documented in the patient hand held record as well as sent to the GP practice for information.

As part of the Shared Care LES, it is expected that ALL patients will have a transfer of care at 3 months, unless there are exceptional circumstances (such as unstable results). We would be grateful if your practice would take over the responsibility for:

- Prescribing the Azathioprine
- Performing the blood tests and monitoring the results (payment via LES)

If patients fail to attend for their monitoring, we would recommend contacting them to arrange one further monitoring appointment but thereafter to stop prescribing their treatment until the monitoring requirements have been met.

The patient carries a hand held monitoring book, which has been kept up to date by the Rheumatology department, and/or GP prescriber, and contains patient information.

Important Information:

- Repeat prescriptions should be retained separately (i.e. highlighted as different to all other repeat prescriptions), so the GP prescriber can ensure monitoring has been undertaken prior to signing and issuing to patient
- Alcohol intake should be limited to nationally recommended levels.
- Azathioprine is relatively safe in pregnancy and when breastfeeding but patients are advised to inform their Rheumatologist if they are planning a pregnancy
- Allopurinol / Febuxostat should not be prescribed (significant interaction)
- Live vaccines should not be given
- Annual flu jab is recommended (to be given by GP practice)
- Avoid exposure to chickenpox and shingles. If infection develops it should be treated aggressively with antiviral medication and Rheumatology dept. can be contacted for advice
- Side effects include: Mild Oral Ulceration /Nausea / Diarrhoea – drug continuation depends on severity and patient wishes.

TPMT (Thiopurine Methyl Transferase) levels are checked before treatment is commenced, in order to predict the 2% of the population most at risk of severe side effects. Treatment is usually started at a dose of 50mg daily and increased by 50mg every 2-4 weeks, to a maximum of 3mg/kg daily according to clinical

response. In practice doses over a total of 200mg daily are seldom necessary. Smaller doses are used in severe renal or hepatic failure and in the elderly.

Monitoring schedule:

- FBC/ U&E / LFT / CRP every 2 weeks until on a stable dose for 6 weeks, then monthly for 3 months
- Then every 3 months unless dose changes
- If dose increase: additional FBC/U&E/LFT at 2, 4 and 6 weeks until on stable dose for 6 weeks then revert to previous schedule
- Results to be entered into hand held monitoring booklet

IF:

WCC	<3.5x 10 ⁹ /l(unless disease related)
Neutrophils	<1.8x 10 ⁹ /l
Platelets	<150x 10 ⁹ /l(unless disease related)
ASTor ALT	> 100

Or Severe Oral Ulceration, Vomiting or Diarrhoea

Stop medication and contact Rheumatology service.

If CRP elevated (>25) and patient symptomatic, inform Rheumatology department. If CRP suddenly elevated without significant change to joint symptoms assess patient for infection. Occasionally patients run a persistently high CRP without joint symptoms – this will usually be mentioned in clinic letters.

Patients should stop Azathioprine if they have significant infection requiring antibiotics (or chickenpox / shingles), and restart once infection treated.

Department Contact details:

Fax: 01709 424276
Telephone Helpline: 01709 424739

Consultants:

Dr Gillian Smith 01709 424275 / 424156
Dr Fiona Fawthrop 01709 424275 / 424156
Dr Rakesh Kumari 01709 424275 / 424156

Nurse Specialists:

Sister Sue Elsey + Sister Louise Hale – Bleep 079 via Switch

Specialist Registrar: available on bleep 101 via Switchboard

AZATHIOPRINE SHARED CARE PROTOCOL

Version:	3 (three)
Ratified by:	ROTHERHAM MEDICINES OPTIMISATION GROUP (RMOG)
Date ratified:	1 st November 2017
Title of originator/author:	Dr Rakesh Kumari, Dr Gillian Smith, Dr Fiona Fawthrop, Rheumatology RFT; Eloise Summerfield, Medicines Management Team RCCG
Title of responsible committee/individual:	Consultants Rheumatology RFT; Medicines Management Team RCCG (BNF 10 Prescribing Advisor)
Date issued:	November 2017
Review date:	November 2020
Target audience:	TRFT consultants & nurses Rotherham CCG GPs & nurses

Version	Date	Author	Status	Comment
Version 1	July 2012	Dr James Maxwell RFT Eloise Summerfield RCCG	Archived	First version onto a two page format
Version 2	July 2014	Dr Gillian Smith, Dr Fiona Fawthrop, RFT; Eloise Summerfield, RCCG	Archived	Review – minimal changes
Version 3	Nov 2017	Dr Rakesh Kumari, Dr Gillian Smith, Dr Fiona Fawthrop, RFT; Eloise Summerfield, RCCG	Current version	Extra blood tests at start & dose change. First paragraph re-worded to reflect current practice. Acceptance letter removed as this is now implied under the Shared Care LES